

News Release

Faye Malek, Family Living Educator
Manitowoc County UW-Extension
P.O. Box 1150, 4319 Expo Drive
Manitowoc, WI 54221-1150

December 11, 2005

Specialized nutrition advice for older adults

Nutrition remains important throughout life. Good nutrition in the later years can help reduce the risk of diseases that strike older adults. This contributes to a higher quality of life, enabling older people to maintain their independence.

In addition to general advice for all Americans, the new *Dietary Guidelines for Americans* have key recommendations for adults over age 50.

- Consume extra Vitamin B-12 from fortified foods or supplements. Almost all multivitamin supplements and fortified foods, such as breakfast cereals, have vitamin B-12 in a form that is very easy for the body to use. As people age, their body's ability to use this vitamin from food tends to decrease, so they need a little extra from fortified food or a supplement.
- Consume plenty of calcium and Vitamin D. As you get older, your bones get thinner and may break more easily. The Dietary Guidelines recommend at least three cups of milk or equivalent foods per day for older adults, along with fortified foods and supplements to boost Vitamin D levels. "n a study at Tufts University, 176 men and 213 women over age 65 reduced their risk of breaking a bone if they took a supplement containing calcium and vitamin D for three years.
- Be physically active. Participating in regular physical activity helps maintain your physical and mental health and quality of life as you get older. Being active may be more difficult as you get older, but even small amounts of physical activity like walking, swimming, or strength training with weights can make a difference in your health. If you have health concerns or a painful condition like arthritis, ask your health care provider to recommend activities that are safe and appropriate for you.
- Reduce sodium and increase potassium. The Dietary Guidelines for Americans recommend that older adults consume no more than 1500 milligrams of sodium and increase their potassium intakes to 4700 milligrams per day. Reducing sodium and increasing potassium are both helpful in preventing high blood pressure. Most of the sodium in our diets comes from salt that is added to processed foods. To reduce sodium, read food labels and look for products with less than five percent of the Daily Value for sodium. Fruits, vegetables and milk are good sources of potassium.
- Play it safe with food. Avoid unpasteurized milk, unpasteurized apple juice, raw sprouts, and raw or undercooked eggs, meat, poultry, fish or shellfish. Deli meats and frankfurters should be reheated until they are steaming hot. Older adults are more vulnerable to foodborne illness, so it is important to take these precautions when selecting and preparing food.

For more information, go online at these sites: Dietary Guidelines for Americans and MyPyramid (<http://www.healthierus.gov> and <http://www.mypyramid.gov>), the National Center for Chronic Disease Prevention and Health Promotion (<http://www.cdc.gov/aging/index.htm>), the U.S. Food and Drug Administration (<http://vm.cfsan.fda.gov/~dms/seniors.html>), and the U.S. Centers for Disease Control and Prevention (http://www.cdc.gov/nccdphp/dnpa/physical/growing_stronger/index.htm and <http://www.cdc.gov/nccdphp/sgr/olderad.htm>).

An EEO/Affirmative Action employer, University of Wisconsin-Extension provides equal opportunities in employment and programming, including Title IX and ADA requirements. Please make requests for reasonable accommodations to ensure equal access to educational programs as early as possible preceding the scheduled program, service or activity.

- END -