

FEEDING DAIRY COWS FOR LONGEVITY

Randy. D. Shaver, Ph.D.

**Professor and Extension Dairy Nutritionist¹
Department of Dairy Science
College of Agricultural and Life Sciences
University of Wisconsin – Madison
University of Wisconsin – Extension**

Take Home Messages

- To increase the longevity of dairy cows the focus needs to be on preventing calving/transition-, mastitis-, reproductive- and lameness-related problems.
- There are a myriad of metabolic and digestive disorders that must be minimized during the transition period through management and feeding in order to prevent early-lactation culls or deaths.
- Transition-cow programs are an on-going challenge for dairy producers and their consultants as many interactions between management and nutrition occur on dairy farms that can influence results. Common sense and cow sense continue to be as important to making these programs work on farms as the research base.
- Dietary Vitamin E and selenium are of major importance for reducing mastitis.
- The primary nutritional factors associated with reproductive performance relate to energy and protein status.
- Laminitis is a multi-factorial disorder that requires detailed emphasis on diet formulation, preparation and delivery, feed bunk management, cow management, and cow comfort to control.
- Supplemental dietary biotin improves hoof health.

¹ Contact at: Room 266 Animal Sciences Building, 1675 Observatory Drive, University of Wisconsin, Madison, WI 53706, (608)263-3491, FAX (608)263-9412, Email: rdshaver@wisc.edu

Introduction

Losses of cows from herds due involuntary (non-selective) culling or death, and the need for the replacement of lost cows, represent a significant income loss for dairy producers. The average removal (reported as either culled or died) was 34% in an analysis of MN-DHIA records for 718 Holstein herds over four years of calvings (Godden et al., 2003). In the 2002 NAHMS survey (USDA, 2002), 76% of all culls were reported as involuntary culls. Udder or mastitis problems, reproductive problems, and lameness/injury comprised 35%, 35%, and 22% of the involuntary culls, respectively (NAHMS, USDA, 2002). Godden et al. (2003) reported that about 25% of the herd removals occurred between 0 and 60 DIM or were likely associated with calving/transition cow issues, and that about 40% of the herd removals occurred after 240 DIM or were likely associated with reproductive failure. A national dairy herd death loss of 4.8% was reported in the 2002 NAHMS survey (USDA, 2002), but death loss averaged 8.1% (range 3.5% to 16.8%) in an analysis of on-farm computerized records from twenty New York herds representing over 12,000 cows (Stone et al., 2005). In the ten herds that recorded their best estimates of cause of death, calving- or transition-related problems were the primary cause for 28% of the deaths and another 10% of the deaths were attributed to mastitis problems (Stone et al., 2005).

The foregoing discussion emphasizes that to increase the longevity of dairy cows the focus needs to be on preventing calving/transition-, mastitis-, reproductive-, and lameness-related culls or deaths. This paper will cover only the nutrition aspects of these areas. It should be noted, however, that management of the cows and their environment is crucial to achieving success in all of these areas and for improving longevity.

Nutrition and Transition-Related Culls

Emerging research and extension guidelines on this topic have been extensively reviewed and discussed in papers published in conference proceedings from around the world for over fifteen years. For a concise summary of this area, refer to the 2001 Dairy NRC (NRC, 2001). Nutrient requirements and diet concentrations needed to meet requirements for far-off and close-up dry cows and fresh cows fed example diets are also provided there (NRC, 2001). Pre-partum intake depression and slow intake ascent postpartum, which are major risk factors in the etiology of metabolic and digestive disorders, make minimizing the incidence of transition-related problems (dystocia, milk fever, ketosis, fatty liver, metritis, displaced abomasum, and sub-acute rumen acidosis) an on-going challenge for dairy producers and their consultants. Body condition at dry-off/calving, dry-period length, length of time on pre-fresh diets, stocking densities in transition pens, abrupt cow movement between facilities, feed delivery, and forage quality are just a few examples of the many interactions between management and

nutrition that can influence the results derived from feeding programs designed for transition cows.

Nutrition and Mastitis-Related Culls

Vitamin E (740 IU supplemental and 1060 IU total) during the dry period reduced the incidence of clinical mastitis by 37% mastitis in the subsequent lactation relative to un-supplemented controls (Smith et al., 1984). Selenium (Se) injection at 21-d before calving in cows fed selenium deficient diets during the dry period did not affect the incidence of clinical mastitis in the subsequent lactation, but reduced the duration of clinical symptoms by 46% relative to un-supplemented controls; this reduction was 62% for the group supplemented with both Vitamin E and selenium (Smith et al., 1984). Weiss et al. (1997) supplemented Vitamin E at either 100 or 1000 IU during the full dry period or 1000 IU followed by 4000 IU during the far-off and close-up dry periods in diets with a low content of selenium. Quarter incidence of clinical mastitis was 25%, 16.7%, and 2.6% during the first 7-d of lactation for cows receiving 100 IU, 1000 IU, or 1000/4000 IU Vitamin E, respectively, during the dry period. Current Dairy NRC recommendations are for 1000 and 500 IU supplemental Vitamin E during the dry and lactation periods, respectively (NRC, 2001). The dietary selenium requirement listed in Dairy NRC (NRC 1989, 2001) and the FDA regulatory limit is 0.3 ppm (DM basis). Other vitamins and trace minerals that play a role in immune function and thus possibly in mastitis prevention include Vitamin A, Beta-carotene, copper, zinc, iron (excess can interfere with copper and zinc absorption), and manganese (NRC, 2001). Dietary Vitamin A and trace mineral recommendations for dry and lactating dairy cows are provided in the 2001 Dairy NRC (NRC, 2001).

Nutrition and Reproductive-Related Culls

Severe negative energy balance, excessive mobilization of body fat, and precipitous loss of body condition in early lactation are associated with poor reproductive performance (Nebel, 2005; Wiltbank et al., 2005). Thus, feeding programs and management practices that promote high energy intake, which is a function of both dietary DM intake and energy density, are critically important especially for transition cows. Apart from the role of fat supplements for increasing diet energy density, specialty fat sources with varying fatty acid profiles aimed at improving reproductive performance are being researched and marketed (Thatcher et al., 2004). Feeding diets too high in protein and (or) too low in rumen-fermentable carbohydrates increases blood- and milk-urea nitrogen and has been associated with poor reproductive performance (Ferguson and Chalupa, 1989). Vitamin E and selenium supplementation during the dry period reduced the incidence of retained placenta, metritis, and cystic ovaries (Harrison et al., 1984), which are disorders that can reduce reproductive performance. Dietary Vitamin A and possibly Beta-carotene and trace minerals play a role in reproductive performance (NRC, 2001). Improved

reproductive performance from diets containing above 0.38% phosphorus (DM basis) is highly unlikely (Wu and Satter, 2000; Tallam et al., 2005).

Nutrition and Lameness-Related Culls

Readers are referred to several reviews and updates on sub-acute rumen acidosis (SARA) and laminitis (Nocek, 1997; Shaver, 2000; Shaver, 2002; Cook et al., 2004; Nordlund et al., 2004; Stone, 2004; Shaver, 2005). SARA is a prevalent problem for dairy herds (Cook et al., 2004; Nordlund et al., 2004) as characterized by having more than 25% of cows sampled via rumenocentesis 4 to 8 hours after a TMR meal with ruminal pH less than 5.5 (Nordlund et al., 2004). Ruminal pH is largely a function of the balance between the production of volatile fatty acids (VFA) from the fermentation of carbohydrates, their neutralization by salivary and dietary buffers, and their removal by absorption across the rumen wall or passage from the rumen (Allen, 1997), and SARA is caused by the consumption of high amounts of ruminally-available carbohydrate, low amounts of effective fiber, or both (Nocek, 1997). Laminitis, an aseptic inflammation of the dermal layers inside the hoof and a major source of lameness for dairy herds, has been linked to SARA (Nocek, 1997; Cook et al., 2004; Nordlund et al., 2004; Stone, 2004).

Despite diet formulation guidelines for neutral detergent fiber (NDF) from forage, total NDF and non-fiber carbohydrate (NFC; NRC, 2001; Stone, 2004), physically-effective NDF (peNDF; Mertens, 1997; Stone, 2004), and starch (Nocek, 1997) along with the feeding of TMR's, some degree of SARA may be inevitable in high-producing dairy herds because chewing time, and as a consequence salivary buffer flow, decline per unit of rumen-fermentable organic matter (RFOM) intake or VFA production as RFOM intake increases (Shaver, 2005). An increased intake of RFOM as milk production increases is a normal consequence of high milk production, because of increases in DM intake and the feeding of higher-concentrate diets to increase dietary energy density (NRC, 2001). Further, bunk management and cow comfort have been implicated as risk factors for SARA and laminitis in dairy herds (Shaver, 2002; Cook et al., 2004; Stone, 2004). The foregoing discussion highlights the challenges that we face in our efforts to minimize SARA and laminitis in today's dairy herds, and underscores that the margin for error in our feeding programs is small.

Ruminal pH declines following meals with the rate of pH decline increasing as meal size increases and as dietary NDF content decreases (Allen, 1997). Dietary supplementation of sodium bicarbonate attenuates the decline in ruminal pH that is observed post feeding (Erdman, 1988), and may attenuate SARA. The recommended inclusion rate for sodium bicarbonate is 0.75 to 1.0% of TMR dry matter. Keunen et al. (2003) reported that cows with experimentally-induced SARA did not attenuate SARA by consuming free-choice sodium bicarbonate.

Researchers reported that 20 mg/cow/day supplemental dietary biotin reduced laminitis-related hoof lesions, white-line separation (Hedges et al., 2001; Fitzgerald et al., 2000; Midla et al., 1998; Potzsch et al., 2003) and sole ulcers (Bergsten et al., 2003), and improved sole ulcer healing (Lischer et al., 2002). Dietary biotin supplementation did not influence ruminal VFA (Zimmerly and Weiss, 2001) or apparent total tract organic matter digestibility (Majee et al., 2003), and it is unlikely that reductions in laminitis-related hoof lesions occur via an attenuation of SARA but rather via biotin's role in keratinization of hoof epidermis (Tomlinson et al., 2004).

Nocek et al. (2000) reported reduced laminitis-related hoof lesions in five commercial dairy herds during the year of dietary supplementation with complexed trace minerals (zinc, manganese, copper and cobalt) relative to the year prior when these herds were not supplemented with this complexed trace mineral mixture. The role of trace minerals in keratinization of hoof epidermis (Tomlinson et al., 2004) could explain this response.

References

- Allen, M. S. 1997. Relationship between fermentation acid production in the rumen and the requirement for physically effective fiber. *J. Dairy Sci.* 80:1447-1462.
- Bergsten, C., P. R. Greenough, J. M. Gay, W. M. Seymour, and C. C. Gay. 2003. Effects of biotin supplementation on performance and claw lesions on a commercial dairy farm. *J. Dairy Sci.* 86:3953-3962.
- Cook, N. B., K. V. Nordlund, and G. R. Oetzel. 2004. Environmental influences on claw horn lesions associated with laminitis and subacute ruminal acidosis in dairy cows. *J. Dairy Sci.* 87:(E. Suppl.):E36-E46.
- Erdman, R. A. 1988. Dietary buffering requirements of the lactating dairy cow: A review. *J. Dairy Sci.* 71:3246-3266.
- Ferguson, J. D., and W. Chalupa. 1989. Impact of protein nutrition on reproduction in dairy cows. *J. Dairy Sci.* 72:746-766.
- Fitzgerald, T., B. W. Norton, R. Elliot, H. Podlich, O. L. Svendsen. 2000. The influence of long-term supplementation with biotin on the prevention of lameness in pasture fed dairy cows. *J. Dairy Sci.* 83:338-344.
- Godden, S. M., S. C. Stewart, R. Cady, S. W. Eicker, J. F. Fetrow, P. Rapnicki, W. Weiland, and H. Spencer. 2003. The relationship between herd rBST-supplementation and other factors and risk for removal for cows in Minnesota Holstein dairy herds. Pages 55-64 *in Proc. Four-State Applied Dairy Nutr. & Mgmt. Conf. MWPS-4SD16.* La Crosse, WI.
- Harrison, J. H., D. D. Hancock, and H. R. Conrad. 1984. Vitamin E and selenium for reproduction of the dairy cow. *J. Dairy Sci.* 67:123-132.
- Hedges, J., R. W. Blowey, A. J. Packington, C. J. O'Callaghan, and L. E. Green. 2001. A longitudinal field trial of the effect of biotin on lameness in dairy cattle. *J. Dairy Sci.* 84:1969-1975.

- Keunen, J. E., J. C. Plazier, I. Kyriazakis, T. F. Duffield, T. M. Widowski, M. I. Lindinger, and B. W. McBride. 2003. Effects of subacute ruminal acidosis on free-choice intake of sodium bicarbonate in lactating dairy cows. *J. Dairy Sci.* 86:954-957.
- Lischer, Ch. J., U. Koller, H. Geyer, Ch. Mulling, J. Schulze, and P. Ossent. 2002. Effect of therapeutic dietary biotin on the healing of uncomplicated sole ulcers in dairy cattle – a double blinded controlled study. *Vet. J.* 163:51-60.
- Majee, D. N., E. C. Schwab, S. J. Bertics, W. M. Seymour, and R. D. Shaver. 2003. Lactation performance by dairy cows fed supplemental biotin and a B-vitamin blend. *J. Dairy Sci.* 86:2106-2112.
- Mertens, D. R. 1997. Creating a system for meeting the fiber requirements of dairy cows. *J. Dairy Sci.* 80:1463-1481.
- Midla, L. T., K. H. Hoblet, W. P. Weiss, M. L. Moeschberger. 1998. Supplemental dietary biotin for prevention of lesions associated with aseptic subclinical laminitis (pododermatitis aseptica diffusa) in primiparous cows. *Am. J. Vet. Res.* 59:733-738.
- National Research Council. 2001. Nutrient Requirements of Dairy Cattle. 7th rev. ed. Natl. Acad. Sci., Washington, DC.
- National Research Council. 1989. Nutrient Requirements of Dairy Cattle. 6th rev. ed. Natl. Acad. Sci., Washington, DC.
- Nebel, R. 2005. Improving dairy cattle reproductive performance. Pages 23-30 *in Proc. Four-State Dairy Nutr. & Mgmt. Conf. MWPS-4SD18.* Dubuque, IA.
- Nocek, J. E., A. B. Johnson, and M. T. Socha. 2000. Digital characteristics in commercial dairy herds fed metal-specific amino acid complexes. *J. Dairy Sci.* 83:1553-1572.
- Nocek, J. E. 1997. Bovine acidosis: Implications on laminitis. *J. Dairy Sci.* 80:1005-1028.
- Nordlund, K. V., N. B. Cook, and G. R. Oetzel. 2004. Investigation strategies for laminitis problem herds. *J. Dairy Sci.* 87:(E. Suppl.):E27-E35.
- Potzsch, C. J., V. J. Collis, R. W. Blowey, A. J. Packington, and L. E. Green. 2003. The impact of parity and duration of biotin supplementation on white line disease lameness in dairy cattle. *J. Dairy Sci.* 86:2577–2582.
- Shaver, R. D. 2005. Feeding to minimize acidosis and laminitis in dairy cattle. Pages 49-60 *in Proc. 67th Cornell Nutr. Conf. for Feed Manufacturers.* East Syracuse, NY.
- Shaver, R. D. 2002. Rumen acidosis in dairy cattle: Bunk management considerations. Pages 75-81 *in Proc. 12th Int. Symp. on Lameness in Ruminants.* Orlando, FL.
- Shaver, R. D. 2000. Feed delivery and bunk management aspects of laminitis in dairy herds fed total mixed rations. Pages 70-77 *in Proc. III Int. Conf. on Bovine Lameness.* Parma, Italy.
- Smith, K. L., J. H. Harrison, D. D. Hancock, D. A. Todhunter, and H. R. Conrad. 1984. Effect of Vitamin E and selenium on incidence of clinical mastitis and duration of clinical symptoms. *J. Dairy Sci.* 67:1293-1300.

- Stone, W. C., C. L. Guard, and D. A. Button. 2005. Why do cows diet? Pages 205-212 *in* Proc. 67th Cornell Nutr. Conf. for Feed Manufacturers. East Syracuse, NY.
- Stone, W. C. 2004. Nutritional approaches to minimize subacute ruminal acidosis and laminitis in dairy cattle. *J. Dairy Sci.* 87:(E. Suppl.):E13-E26.
- Tallam, S. K., A. D. Ealy, K. A. Bryan, and Z. Wu. 2005. Ovarian Activity and Reproductive Performance of Dairy Cows Fed Different Amounts of Phosphorus. *J Dairy Sci.* 88:3609 - 3618.
- Thatcher, W. W., C. R. Staples, J. Santos, and S. Juchem. 2004. Effects of lipids on reproductive responses in lactating dairy cattle. Pages 103-129 *in* Proc. Midwest Dairy herd Health Conf. Eau Claire, WI.
- Tomlinson, D. J., Mulling, C. H., and T. M. Fakler. 2004. Formation of keratins in the bovine claw: Roles of hormones, minerals, and vitamins in functional claw integrity. *J. Dairy Sci.* 87:797-809.
- USDA, National Animal Health Monitoring System. Dairy 2002. Part 1: Reference of Dairy Health and Management in the United States.
- Weiss, W. P., J. S. Hogan, D. A. Todhunter, and K. L. Smith. 1997. Effect of Vitamin E in diets with a low concentration of selenium on mammary gland health of dairy cows. *J. Dairy Sci.* 80:1728-1737.
- Wiltbank, M., H. Lopez, R. Sartori, and A. Gument. 2005. Current management of positive and negative reproductive impacts of high energy intake. Pages 53-64 *in* Proc. Four-State Dairy Nutr. & Mgmt. Conf. MWPS-4SD18. Dubuque, IA.
- Wu, Z., and L. D. Satter. 2000. Milk Production and Reproductive Performance of Dairy Cows Fed Two Concentrations of Phosphorus for Two Years. *J. Dairy Sci.* 83:1052-1063.
- Zimmerly, C. A. and W. P. Weiss. 2001. Effects of supplemental dietary biotin on performance of Holstein cows during early lactation. *J. Dairy Sci.* 84:498-506.