

## Bean Leaf Beetle

*Karen Delahaut, UW-Madison IPM Program*

Bean leaf beetles (*Cerotoma trifurcate*) are becoming more of a problem in Wisconsin than they have been in the past. Warmer winters may allow overwintering adults to survive in record numbers. Host plants include snap and lima beans, peas, soybeans, and soybeans as well as leguminous weeds.



### Appearance

Adult bean leaf beetles are ¼ inch long and range in color from yellow-green to red. They have 4 black spots on their backs, a black triangle on the wing covers behind the head, and a black band surrounding the outer margins of their wings. The larvae are slender and white with dark areas on both ends.

### Symptoms and Effects

Adult bean leaf beetles chew small holes into the leaves while feeding from the lower leaf surface. Adult feeding on young plants can reduce vigor and

yield. They can also feed on the developing pods causing distortion as well as creating a point of entry for disease organisms. The larvae feed on the roots and nodules as well as the stems below the soil line and may girdle small plants.

### Life Cycle

The beetles overwinter as adults in leaf litter, plant debris, weeds, and grassy fields near fields that had beans planted in them the previous year. They become active in mid-May to early June when the weather warms. Before the host crop emerges, bean leaf beetles feed on alfalfa, clover, and prairie legumes. The beetles often migrate into vegetable crops when nearby alfalfa is cut. After mating, females lay small clusters of lemon-shaped, orange eggs in the soil near the base of the plant. The eggs hatch in 1-3 weeks and the larvae immediately begin feeding on the roots and underground stems. The larvae feed for 30 days before pupating. First generation adults appear in mid-July. A second generation of adults appears in late August and September. These will become the overwintering adults that will emerge next spring.





## Scouting Suggestions

Preliminary thresholds established by the University of Minnesota indicate that beans should be treated when there are 6-10 shot holes per leaf on 10% of the plants when plants are between the seedling stage and when they produce their first trifoliolate leaves.

## Control

**Cultural** Bean leaf beetles are very difficult to control without chemicals. Because of the highly mobile nature of the beetles, control options are limited. Discourage overwintering beetles by destroying plant debris and keeping vegetation mowed near susceptible crops. Beans planted in early to mid-June are more likely to escape damage by the overwintering adults. Floating row covers placed over the crop at planting in small gardens will exclude the beetles from getting to the crop but this isn't practical in large plantings. Be sure to firmly secure all edges of the row cover so the beetles don't crawl beneath.

**Chemical** Commercial snap bean growers who have had problems with bean leaf beetles in the past should use an imidicloprid insecticidal seed treatment at planting. Insecticides used to control other insects such as the European corn borer will also control bean leaf beetles. Organic growers can use Rotenone, pyrethrum or neem and those not growing organically can also use carbaryl and permethrin. It's important to treat early and get thorough coverage of the foliage particularly the lower leaf surface. Refer to UWEX publication A3422 "Commercial Vegetable Production in Wisconsin" for a list of recommended products.

**For pesticide recommendations:** See UW-Extension Bulletin A3422 or contact your County Extension Agent.

© 2003 by the Board of Regents of the University of Wisconsin System doing business as the division of Cooperative Extension of the University of Wisconsin Extension.

An EEO/Affirmative Action employer, University of Wisconsin Extension provides equal opportunities in employment and programming, including Title IX and ADA requirements.

References to pesticide products in this publication are for your convenience and are not an endorsement or criticism of one product over similar products. You are responsible for using pesticides according to the manufacturer's current label directions. Follow directions exactly to protect the environment and people from pesticide exposure. Failure to do so violates the law.

Thanks to Phil Pellitteri & Lis Friemoth for reviewing this document.