

## Management Practices and Reported Antimicrobial Usage on Conventional and Organic Dairy Farms<sup>1</sup>

A. G. Zwald,<sup>1</sup> P. L. Ruegg,<sup>1</sup> J. B. Kaneene,<sup>2</sup> L. D. Warnick,<sup>3</sup>  
S. J. Wells,<sup>4</sup> C. Fossler,<sup>4</sup> and L. W. Halbert<sup>2</sup>

<sup>1</sup>Department of Dairy Science  
University of Wisconsin  
Madison 53706

<sup>2</sup>Population Medicine Center  
College of Veterinary Medicine  
Michigan State University  
East Lansing 48824

<sup>3</sup>Department of Population Medicine and Diagnostic Sciences  
College of Veterinary Medicine  
Cornell University  
Ithaca, NY 14850

<sup>4</sup>Department of Clinical and Population Science  
College of Veterinary Medicine  
University of Minnesota  
St. Paul 55108

### ABSTRACT

The primary objective was to compare reported antimicrobial usage between conventional and organic dairy farms. A secondary objective was to contrast selected management characteristics of conventional and organic dairy herds. A questionnaire was administered on site to selected dairy farmers located in Michigan, Minnesota, New York, and Wisconsin. Organic herds ( $n = 32$ ) were smaller and produced less milk than conventional herds ( $n = 99$ ). Lactating cows in organic dairies were more likely to be housed in tie stalls, whereas most conventional dairies housed cows in free stalls and milked in a parlor. Total mixed rations and purchased feeds were used on more conventional dairy farms compared with organic dairy farms. Conventional dairy producers were more likely to use advice from veterinarians for recommendations of treatment, and organic dairy producers were more likely to rely on advice from other farmers. Based on recall of antibiotic usage in the previous 60 d, 5.1, 84.9, 9.1, and 0.9% of farmers with conventional herds reported treatment of none, 1 to 10%, 11 to 25%, and >25% of milk cows, respectively. Most organic farmers (90.6%) reported no antibiotic treatments of milk cows, whereas 9.4% reported treating 1 to 10% of milk cows. Ceftiofur was the most commonly reported antibiotic for both farm

types. Milk replacer containing antibiotics was reportedly used on 49.5% of conventional herds but only on one organic herd (3.1%). Antibiotics were used in heifer calves on 74.7% of conventional herds versus 21.9% of organic herds. Antibiotics to treat mastitis were used on 79.8% of conventional herds but on none of the organic herds. Most organic farms were in compliance with standards in advance of implementation of regulations. (**Key words:** antibiotics, antimicrobials, organic, management)

**Abbreviation key:** CON = conventional, ORG = organic.

### INTRODUCTION

Antibiotic treatment of dairy cows for infectious diseases is a relatively common and necessary occurrence (McEwen et al., 1991). The overall availability and use of antimicrobials have been estimated in a US population-based survey (National Animal Health Monitoring System, 1997); however, there are few data available on the use of individual antimicrobials. It is difficult to measure antimicrobial usage on farms because of the difficulty of obtaining an accurate assessment of dosage and duration of treatment. Antibiotics are administered to animals through injections (e.g., intramuscular, intravenous, or subcutaneous), orally, topically, or via intramammary or intrauterine infusion. A survey of US veterinarians found that the drugs most often used or prescribed were antibiotics, followed by antiinflammatories and tranquilizers or analgesics (Sundlof et al., 1995).

---

Received April 3, 2003.

Accepted July 16, 2003.

Corresponding author: P. L. Ruegg; e-mail: plruegg@wisc.edu.

<sup>1</sup>Supported by UDSA-NRI Program on Epidemiological Approaches to Food Safety.

Several types of antimicrobials are commonly used in food animals (Mitchell et al., 1998). Antimicrobial classes include beta-lactams (e.g., penicillin, ampicillin, and cephalosporin), tetracyclines (e.g., oxytetracycline, tetracycline, and chlortetracycline), aminoglycosides (e.g., streptomycin, neomycin, and gentamycin), macrolides (e.g., erythromycin), lincosamides (e.g. lincomycin and pirlimycin), and sulfonamides (e.g., sulfamethazine and others) (Mitchell et al., 1998; Hoeben et al., 1998). A survey of US veterinarians reported that antibiotics were the drugs most often used to treat lactating dairy cows; the most common reason for treatment was mastitis therapy (Mitchell et al., 1998).

The use of antimicrobials in agriculture is not universally accepted, and the market for organic agricultural products has increased. Although the organic market has been growing at 20 to 30% per year in the United States, scientific literature concerning organic dairy farming is relatively rare (Green, 2000). As of October 2002, the National Organic Program, under the Agricultural Marketing Service and USDA, defined standards for organic production and handling. The national organic standards address the methods, practices, and substances used in producing and handling crops, livestock, and processed agricultural products. The requirements apply to the production process, not to measurable properties of the product itself. It is a requirement that livestock destined for slaughter are raised under organic management from the last third of gestation. During transition to organic status, cattle intended for dairy production do not need to be managed organically until a year before commencing lactation. After attaining organic status, all animals on the dairy must be raised under organic rules from the last third of gestation. Animals sold as organic may not have received hormones to promote growth, or antibiotics for any reason. Preventive management practices, including vaccines, are allowed. Producers are prohibited from withholding treatment from a sick or injured animal; however, products derived from animals treated with a prohibited medication may not be sold as organic. All organically raised cattle must have access to the outdoors. The current national organic standards were implemented after this study ended. There are no studies contrasting the use of antimicrobials based on specific farming systems (organic vs. conventional). The primary objective of this study was to compare reported antimicrobial usage between conventional and organic dairy herds. A secondary objective was to contrast selected management characteristics of the conventional and organic dairy herds enrolled in this study.

## MATERIALS AND METHODS

### Recruitment and Herd Selection

Study farms (n = 131) were recruited from dairy herds located in the states of Michigan, Minnesota, New York,

and Wisconsin between May 2000 and March 2001. Recruitment methods differed slightly among states and between farm types [conventional (**CON**) and organic (**ORG**)]. Conventional herds were identified by obtaining a list of dairy farms licensed to ship milk from the department of agriculture of each respective state. Organic herds were required to be certified by an organic certifying agency for at least 3 yr. There is no central database of ORG herds, so lists of ORG farms were obtained through personal contacts, organic milk cooperatives, and independent organic certifying organizations. In some states, all known ORG herds within each state were contacted by phone to determine eligibility and interest in participation in this study. In Minnesota and Wisconsin, letters were sent to ORG herds to solicit participation in this study.

Conventional herds were contacted by letters (500 to 571 per state for a total of 2102 letters) mailed to randomly selected herds located in counties near each university. Nonresponders received multiple mailings in order to increase participation rate. The letters included a postage paid return postcard. Farms that indicated willingness to participate in the study (n = 295) were screened to ensure that they met predefined criteria for eligibility: preweaned calves and heifers raised on the farm property, a herd record system, unique identification for all cows and calves, >90% of the herd population was Holstein, the farm shipped milk year round, and the farm owner agreed to participate in the study. The absence of any single eligibility criterion resulted in exclusion from the study. The four herd size categories of 30 to 49 cows (n = 21), 50 to 99 cows (n = 42), 100 to 199 cows (n = 33), and 200 or more cows (n = 35) were filled by random selection from eligible herds.

### Questionnaire

A 64-question survey instrument (available on request) including inventory and herd size (3 questions), herd expansion status (2 questions), housing (5 questions), feed and water system (10 questions), calf management and feeding (12 questions), production and health (10 questions), manure management (9 questions), and antimicrobial use (13 questions) was administered. The questionnaire was adapted from a previously published survey (National Animal Health Monitoring System, 1997). For many questions, producers could select all options that applied specifically to their farms. An individual animal area was defined as an area where animals could not commingle with other dairy animals. A multiple animal area was defined as an area where animals did have opportunities for direct contact with other dairy animals. Diets of dry cows were recorded separately from diets fed to lactating cows. An

additional 10 questions (addressing the separation and sale of animals that received antimicrobials and type of products used for treatments) were included on the questionnaire administered to ORG dairy farms. To reduce recall bias, all questions referred to practices used in the previous 60 d (unless otherwise noted). For one question, we asked study participants to estimate the percentage of cattle that had received at least one antibiotic injection or oral dose of antibiotics within the previous 60 d for adult cows (milking or dry), bred heifers, and heifer calves (preweaned or weaned). This question did not include the use of intramammary products or topical administration of antibiotics but did include injectable and oral treatments given by farm workers or veterinarians. A separate question addressed the use of intramammary dry cow therapy. The questionnaire was pretested by administration to several farmers and 2 faculty members located in Minnesota. The questionnaire took approximately 45 min to 2 h to complete and was administered during a farm visit. Data were collected during a meeting with the farmer or herdsman and included an interview, review of product labels (i.e., milk replacer and antibiotics), and information obtained during a farm tour.

### Statistical Analyses

Data from all states were entered into a central database for analysis. Data were stratified by herd size ( $\leq 120$  total cattle = small,  $>120$  total cattle = large) for initial data analysis. There was no significant difference in the number of cows between ORG and CON herds for small herds (89.9 cows [CON], 84.0 cows [ORG];  $P = 0.41$ ). Large CON herds contained more cattle compared with large ORG herds (398.0 cows [CON], 262.9 cows [ORG];  $P = 0.02$ ). The data were stratified by herd size (small or large) with herd type (ORG or CON) forming the rows and the response variable (yes or no for specified antibiotics) forming the columns and were analyzed using the Cochran-Mantel-Haenszel option in PROC FREQ (SAS, 1999). Herd size was not associated with antibiotic usage so data were combined for further analysis. The PROC FREQ chi-square analysis and Fisher's exact test (SAS, 1999) were used to identify associations between herd type and the use of selected antibiotics or other treatments. Continuous variables were analyzed using PROC TTEST and PROC MEANS (SAS, 1999).

## RESULTS

### Herd Characteristics

Herd enrollment was distributed among the participating states: 30 (CON) and 2 (ORG); 23 (CON) and 10

(ORG); 25 (CON) and 9 (ORG); and 21 (CON) and 11 (ORG) for Michigan, Minnesota, New York, and Wisconsin, respectively. The distribution of enrollment by herd type was: 12 (ORG) and 9 (CON); 12 (ORG) and 30 (CON); 4 (ORG) and 29 (CON); and 4 (ORG) and 31 (CON) for herd size categories of 30 to 49, 50 to 99, 100 to 199, and  $>200$  total cows, respectively.

Organic herds were smaller and produced less milk than CON herds (Table 1). One ORG herd was allowed to remain in the study even though it contained only 26 adult dairy cows on the day of the herd visit. Higher SCC values were reported for proportionally more ORG herds compared with SCC values reported by CON herds (Table 1). A larger proportion of cows in CON herds were milked in a pit parlor, whereas a larger proportion of cows in ORG herds were milked in a tie stall or stanchion barn (Table 1).

### Housing and Management

Preweaned calves in ORG herds were housed in areas that permitted less contact with other animals, but there was no significant association between herd type and specific housing types such as the housing of preweaned calves nor with the housing of weaned calves (Table 2). However, significantly more CON herds housed lactating cows in free stalls and had maternity housing separate from lactating cows compared with ORG herds (Table 2). Calves in more CON herds were fed individually (i.e., no access to group feeding such as shared troughs) compared with ORG herds. The use of separate facilities to house sick animals separately from lactating cows and the washing of calf pens were not influenced by herd type.

Nutritional management was affected by herd type. A significantly greater percentage of CON herds fed lactating cows a TMR (79.8 vs. 31.3%,  $P < 0.001$ ), a transition ration (71.7 vs. 31.3%,  $P < 0.001$ ), and anionic salts (33.3 vs. 3.1%,  $P = 0.001$ ) to close-up cows. Whole soybeans and soybean meal were the only purchased feeds reported on ORG dairies, but significantly less ORG (46.9%) compared with CON (90.9%) herds used them ( $P < 0.001$ ). The use of whole cottonseed or hulls, brewer's byproducts, and tallow or animal fat was not reported for ORG dairies, whereas these products were used on 50.5, 48.5, and 60.6% of CON farms, respectively ( $P < 0.001$ ). No organic farmers reported the use of meat and bone meal or blood meal, whereas these products were used on 27.3% of CON herds ( $P < 0.02$ ).

Individual automatic water bowls were used on significantly more ORG herds (Table 3). Occasional use of surface water by milking cows and primary use of surface water by dry cows occurred in significantly more ORG herds (Table 3). Chlorinated drinking water was

**Table 1.** Selected descriptive statistics for herds by herd type.

	Conventional					Organic				
	N	Mean	SE	Min	Max	N	Mean	SE	Min	Max
Total adult cows <sup>1</sup>	99	192.2 <sup>a</sup>	17.3	30.0	875.0	32	90.5 <sup>b</sup>	16.8	26.0	368.0
Total cattle <sup>2</sup>	99	348.2 <sup>a</sup>	30.0	50.0	1693.0	32	173.4 <sup>b</sup>	28.2	47.0	677.0
Rolling herd average (kg)	87	9896.9 <sup>a</sup>	142.9	6905.9	13,063.5	22	7555.9 <sup>b</sup>	235.7	5443.1	9911.0
Daily milk yield per cow (kg)	28	30.9 <sup>a</sup>	1.1	20.4	39.0	14	22.8 <sup>b</sup>	1.2	13.6	30.8
SCC <sup>3</sup> (cells per ml)		(%)					(%)			
<100,000	1	(1.0)				1	(3.1)			
100,000–199,999	30	(30.3)				2	(6.3)			
200,000–299,999	42	(42.4)				14	(43.8)			
300,000–399,999	18	(18.2)				10	(31.3)			
400,000–499,999	6	(6.1)				5	(15.6)			
≥500,000	2	(2.0)				0	(0.0)			
Milking facility <sup>4</sup>										
Pit parlor	60	(60.6)				6	(18.8)			
Flat parlor	7	(7.1)				0	(0.0)			
Tie stall or stanchion	32	(32.3)				26	(81.3)			

<sup>a,b</sup>Means within the same row with different superscripts differ significantly,  $P < 0.002$ .

<sup>1</sup>Includes lactating and dry cows.

<sup>2</sup>Includes adults and youngstock.

<sup>3</sup>SCC category was associated with herd type,  $P = 0.04$ .

<sup>4</sup>Type of milking facility was associated with herd type,  $P < 0.001$ .

**Table 2.** Cattle housing for conventional (n = 99) and organic (n = 32) dairy herds.

	Conventional		Organic		$P^1$
	Number of herds	(%)	Number of herds	(%)	
Housing of preweaned calves <sup>3</sup>					
Hutch	56	(56.6)	13	(40.6)	NS <sup>2</sup>
Tie stall	2	(2.0)	2	(6.3)	NS
Tied in stanchion barn	10	(10.1)	4	(12.5)	NS
No contact with other animals	13	(13.1)	20	(62.5)	<0.001
Contact with other animals possible	43	(43.4)	9	(28.1)	0.12
Housing of weaned calves <sup>3</sup>					
Hutch	4	(4.0)	0	(0.0)	NS
Free stall	22	(22.2)	7	(21.9)	NS
Tie stall	6	(6.1)	3	(9.4)	NS
Contact with other animals possible	89	(89.9)	30	(93.8)	NS
Housing of lactating cows <sup>3</sup>					
Free stall	73	(73.7)	11	(34.4)	<0.001
Tie stall	35	(35.4)	25	(78.1)	<0.001
Contact with other animals possible	14	(14.1)	11	(34.4)	0.011
Maternity housing <sup>3</sup>					
Free stalls	6	(6.1)	2	(6.3)	NS
Tie stalls	8	(8.1)	9	(28.1)	<0.01
No contact with other animals	43	(43.4)	16	(50.0)	NS
Contact with other animals possible	58	(58.6)	13	(40.6)	0.08
Separate from lactating cows	82	(82.8)	16	(50.0)	<0.001
Sick cows housed separate from lactating cows	41	(41.4)	10	(31.3)	NS
Preweaned calves fed on an individual basis	92	(92.9)	25	(78.1)	<0.05
Calf pens washed on a regular basis	51	(51.5)	17	(53.1)	NS
No calf pens or hutches used to house calves	11	(11.1)	10	(31.3)	<0.05

<sup>1</sup>Contrast between conventional and organic.

<sup>2</sup>NS =  $P > 0.15$ .

<sup>3</sup>Column totals within herd type can exceed number of enrolled herds because housing options were not exclusive choices.

**Table 3.** Water sources by herd type for conventional (n = 99) and organic (n = 32) dairy herds.

	Conventional		Organic		P
	Number of herds	(%)	Number of herds	(%)	
Milking cows <sup>1</sup>					
Individual automatic water bowl	36	(36.4)	26	(81.3)	<0.001
Automatic water bowl shared by group	37	(37.4)	13	(40.6)	NS <sup>2</sup>
Water tank for multiple cows	83	(83.8)	26	(81.3)	NS
Occasional access to surface water	10	(10.1)	10	(31.3)	<0.01
Main water source is surface water	3	(3.0)	3	(9.4)	0.14
Dry cows <sup>1</sup>					
Individual automatic water bowl	21	(21.2)	15	(46.9)	<0.01
Automatic water bowl shared by group	47	(47.5)	17	(53.1)	NS
Water tank for multiple cows	61	(61.6)	19	(59.4)	NS
Occasional access to surface water	14	(14.1)	8	(25.0)	0.15
Main water source is surface water	9	(9.1)	7	(21.9)	0.06
Source of drinking water <sup>1</sup>					
Well	96	(97.0)	32	(100.0)	NS
Municipal water	1	(1.0)	0	(0.0)	NS
Surface water	24	(24.3)	9	(28.1)	NS
Drinking water for cows chlorinated	13	(13.1)	0	(0.0)	<0.05

<sup>1</sup>Column totals within herd type can exceed number of enrolled herds because options were not exclusive choices.

<sup>2</sup>NS = P > 0.15.

used on significantly more CON dairies. There was no significant association of herd type with group access to individual automatic water bowls, the use of water tanks for multiple cows, occasional use of surface water by dry cows, and use of surface water as main water sources for milking cows. Similarly, there was no significant association between water source (well water, municipal source, or surface water) and herd type.

Physical contact between poultry and dairy cattle was reported on significantly more ORG herds (37.5%) compared with CON herds (11.1%;  $P = 0.001$ ). Herd type was not associated with physical contact with beef cattle, horses, pigs, sheep, goats, dogs, cats, or wild geese ( $P > 0.20$ ).

A treated cow was defined as a cow that had received antibiotics that required a withholding period that had not yet ended. Whole milk obtained from treated cows was fed to calves on significantly more CON herds (38.4% [CON] vs. 0.0% [ORG],  $P < 0.001$ ), whereas whole milk obtained from untreated cows was fed to calves on more ORG herds (25.3% [CON] vs. 93.8% [ORG],  $P < 0.001$ ). The majority of farms fed  $\geq 3.78$  L of colostrum to newborn calves (80.0% [CON] vs. 63.4% [ORG],  $P = 0.29$ ).

Diagnosis of Johne's disease was more commonly reported on CON farms (48.5%) compared with ORG farms (25.0%;  $P = 0.02$ ). There was no significant association between recall of previous diagnosis of *Salmonella* or bovine viral diarrhea and herd type ( $P > 0.11$ ). The use of coliform mastitis vaccines was more common on CON (47.5%) compared with ORG dairy farms (15.6%;  $P = 0.001$ ).

## Antimicrobial Usage

**Records and recommendations for antibiotic use.** Antibiotic treatment of both lactating and nonlactating cows was recorded on significantly more CON herds (Table 4). Likewise, reliance on veterinarian recommendations for antibiotic use, dosage, and withdrawal time was more common on CON herds (Table 4). Reliance on personal experience for recommendations for antibiotic use, dosage, and withdrawal time was cited by a high percentage of farmers on both ORG and CON herds. Advice obtained from other farmers for antibiotic withdrawal time was utilized by a low percentage of farmers on both CON and ORG herds. The product label was more frequently chosen as a source of withdrawal time information on CON, compared with ORG, herds. Recommendations obtained from other farmers for antibiotic use and dosage were more frequently cited by ORG farmers than on CON farmers. Antibiotic treatments of calves and heifers were recorded on approximately one-third of farms, regardless of herd type.

**General antibiotic usage.** Based on recall of the previous 60 d, there was a significant association between the percentage of calves, bred heifers, and adult cows that received at least one antibiotic dose in the past 60 d and herd type (Table 5). One to ten percent of cows received at least one dose of antibiotics on the majority of CON herds, whereas no cows received antibiotics on the majority of ORG herds. More use of all antibiotics included in the survey was reported on CON herds compared with ORG herds (Table 6).

**Table 4.** Antibiotic recommendations by herd type for conventional (n = 99) and organic (n = 32) dairy herds.

	Conventional		Organic		P
	Number of herds <sup>1</sup>	(%)	Number of herds <sup>1</sup>	(%)	
Antibiotic treatment records					
Lactating cows	71	(71.7)	6	(18.8)	<0.001
Nonlactating cows	58	(58.6)	7	(21.9)	<0.001
Calves and heifers	36	(36.4)	11	(34.4)	NS <sup>2</sup>
Recommendations for antibiotic use from					
Veterinarian	99	(100.0)	23	(71.9)	<0.001
Pharmaceutical representative	12	(12.1)	5	(15.6)	NS
Personal experience	75	(75.8)	27	(84.4)	NS
Product label	39	(39.4)	10	(31.3)	NS
Other farmers	27	(27.3)	22	(68.8)	<0.001
Recommendations for antibiotic dosage from					
Veterinarian	96	(97.0)	22	(68.8)	<0.001
Pharmaceutical representative	8	(8.08)	3	(9.4)	NS
Personal experience	65	(65.7)	25	(78.1)	NS
Product label	55	(55.7)	14	(43.8)	NS
Other farmers	10	(10.1)	18	(56.3)	<0.001
Recommendations for antibiotic withdrawal time from					
Veterinarian	91	(91.9)	12	(37.5)	<0.001
Pharmaceutical representative	4	(4.0)	0	(0.0)	NS
Personal experience	54	(54.6)	8	(25.0)	<0.01
Product label	66	(66.7)	4	(12.5)	<0.001
Other farmers	9	(9.1)	5	(15.6)	NS

<sup>1</sup>Column totals within herd type can exceed number of enrolled herds because options were not exclusive choices.

<sup>2</sup>NS =  $P > 0.15$ .

Antibiotics were used to treat dairy cattle on 15 ORG herds (50%). Of the 15 herds using antibiotics, 14 (93.3%) reported that they separated dairy cattle that received antibiotics from the rest of the ORG herd. The term "separated" referred to selling the animal or moving her to a location that did not allow physical contact with the rest of the ORG herd. The separation was permanent for 13 (92.9%) of the 14 herds. Treated animals were used for ORG milk production after a withdrawal period had passed on 3 ORG farms (20%). However, no animals previously treated with antibiotics were sold for ORG meat production.

**Antibiotic usage in replacement animals.** Antibiotics were used to treat respiratory disease and diarrhea in calves on significantly more CON herds compared with ORG herds (Table 7). The use of milk replacer containing antibiotics was reported on nearly

half of the CON herds (49.5%) compared with only one ORG herd (3.1%) ( $P < 0.001$ ). There was no difference in the use of calf starter based on herd type (87.9% [CON], 78.2% [ORG],  $P = 0.14$ ) There was no significant association between herd type and the use of calf starter without antibiotics (63.6% [CON], 68.8% [ORG];  $P = 0.60$ ), or use of calf starter with antibiotics (24.3% [CON], 9.4% [ORG],  $P = 0.07$ ). Oxytetracycline (46.5%) and neomycin (42.4%) were the antibiotics most commonly reported used in milk replacers or calf starters on CON herds. The use of medications other than coccidiostats and ionophores in the feed or water of weaned calves and heifers was reported only by CON farmers (17.3%;  $P = 0.01$ ), but only 4.0% of the farmers reported using these medications continuously. Chlortetracycline or another antibiotic was used as a medication in

**Table 5.** Use of antibiotics in animals on organic and conventional dairy herds in the 60 d preceding the interview.<sup>1</sup>

Percent of herd receiving at least 1 dose of antibiotics	Heifer calves <sup>1</sup>		Bred heifers		Adult cows <sup>1</sup>	
	Conventional	Organic	Conventional	Organic	Conventional	Organic
Zero	25.3%	78.1%	82.8%	0.0%	5.1%	90.6%
1–10%	57.6%	21.9%	17.2%	0.0%	84.9%	9.4%
11–25%	10.1%	0.0%	0.0%	0.0%	9.1%	0.0%
>25%	7.0%	0.0%	0.0%	0.0%	0.9%	0.0%

<sup>1</sup>Herd type was significantly associated with percentage of herd receiving antibiotics,  $P < 0.001$ .

**Table 6.** Types of antibiotics reported used in previous 60 d by organic and conventional dairy herds.

Antibiotic	Conventional	Organic	<i>P</i>
Penicillin type	85.9%	6.5%	<0.001
Cephalosporin type	77.85	9.7%	<0.001
Tetracycline type	41.4%	3.2%	<0.001
Sulfonamides	27.3%	0.0%	<0.001
Florfenicol	26.3%	6.5%	<0.05
Other antibiotics	37.4%	9.7%	<0.01

the feed or water of weaned calves and heifers on 10.1% of CON herds.

**Coccidiostats and ionophores.** Use of decoquinatate in rations of both preweaned calves (60.6% [CON], 12.5% [ORG],  $P < 0.001$ ), and weaned calves (24.3% [CON], 3.1% [ORG],  $P = 0.004$ ) was reported in more CON herds compared with ORG herds. Similarly, more CON herds reported the use of lasalocid to feed preweaned calves (39.4% [CON], 3.1% [ORG],  $P < 0.001$ ). There was no significant difference between CON and ORG herds for use of amprolium in preweaned calves (10.1% [CON], 3.1% [ORG],  $P = 0.22$ ), or the use of other coccidiostats and ionophores in weaned calves (4.0% [CON], 6.3% [ORG],  $P = 0.60$ ). The use of coccidiostats and ionophores in bred heifers was not reported for ORG herds. Monensin (22.2%;  $P = 0.002$ ) and decoquinatate (5.05%;  $P > 0.11$ ) were reported to be used on CON herds. Monensin was not used on any ORG herds but was used in weaned calves on 25.3% of CON herds ( $P = 0.001$ ) and in preweaned calves on 9.1% of CON herds ( $P > 0.11$ ).

**Antibiotic usage in adult cows.** Respiratory disease, mastitis, metritis or retained placenta, and foot

**Table 8.** Use of systemic antibiotics to treat selected diseases in adult cows.

Disease	Conventional	Organic	<i>P</i>
Respiratory disease	97.0%	12.5%	<0.01
Mastitis	79.8%	0.0%	<0.001
Metritis or retained placenta	79.8%	3.1%	<0.001
Foot problems	82.8%	6.3%	<0.001

problems were treated more commonly with systemic antibiotics on CON herds compared with ORG herds (Table 8).

Ceftiofur was used in cows to treat respiratory disease, retained placenta or metritis, and foot problems on some ORG dairies (Table 9). More CON farmers reported the use of ceftiofur, tetracycline, penicillin, ampicillin, and sulfonamides to treat respiratory disease in cows compared with ORG farmers ( $P < 0.004$ ), yet overall use of many of these antibiotics was relatively low (Table 9). There was no significant difference between herd types for use of florfenicol, tilmicosin, and other antibiotics for respiratory disease in cows because of the low use of these antibiotics in CON herds and the absence of use in ORG herds ( $P > 0.07$ ). The use of systemic antibiotics to treat mastitis was not reported in ORG herds, but a variety of medications were reported to treat mastitis in CON herds, and additional products in the “other” category were reported in about one-fourth of CON herds ( $P < 0.007$ ) (Table 9). There was no significant difference between herd type for systemic use of ceftiofur, sulfonamides, and erythromycin for mastitis in cows due to the low use of these antibiotics in CON herds and absence of use in ORG herds ( $P > 0.11$ ). The use of ceftiofur to treat retained placenta

**Table 7.** Antibiotic usage in calves for conventional (n = 99) and organic (n = 32) dairy herds.

Antibiotic	Respiratory disease					Calf diarrhea				
	Conventional		Organic		<i>P</i>	Conventional		Organic		<i>P</i>
	N	(%)	N	(%)		N	(%)	N	(%)	
Use antibiotics to treat	96	(97.0)	11	(34.4)	<0.001	78	(78.8)	7	(21.9)	<0.001
Ceftiofur <sup>1</sup>	37	(37.4)	0	(0.0)	<0.001	11	(11.1)	0	(0.0)	0.06
Tetracyclines <sup>1</sup>	30	(30.3)	1	(3.1)	<0.01	21	(21.2)	0	(0.0)	<0.01
Penicillin <sup>1</sup>	36	(36.4)	5	(15.6)	<0.05	9	(9.1)	9	(9.4)	NS <sup>2</sup>
Ampicillin <sup>1</sup>	8	(8.1)	0	(0.0)	NS					
Lincomycin-spectinomycin <sup>3</sup>	4	(4.0)	1	(3.1)	NS					
Tilmicosin <sup>1</sup>	45	(45.5)	3	(9.4)	<0.01					
Florfenicol <sup>1</sup>	36	(36.4)	2	(6.3)	0.001	13	(13.1)	1	(3.1)	0.11
Gentamycin <sup>3</sup>	2	(2.0)	0	(0.0)	NS					
Trimethoprim-sulfa <sup>3</sup>						26	(26.3)	1	(3.1)	<0.01
Spectinomycin <sup>3</sup>						11	(11.1)	0	(0.0)	0.06
Enrofloxacin <sup>3</sup>	3	(3.0)	0	(0.0)	NS					
Other	16	(16.2)	2	(6.3)	NS	22	(22.1)	2	(6.3)	<0.05

<sup>1</sup>Antibiotic specifically mentioned in question.

<sup>2</sup>NS =  $P > 0.15$ .

<sup>3</sup>Antibiotic volunteered by producers in the “other” category.

**Table 9.** Antibiotic usage in adult cows in conventional (n = 99) and organic (n = 32) herds for selected ailments (percent).<sup>1,2</sup>

Antibiotic	Respiratory disease				Mastitis				Retained placenta or metritis				Foot problems			
	Conventional		Organic		Conventional		Organic		Conventional		Organic		Conventional		Organic	
	N	(%)	N	(%)	N	(%)	N	(%)	N	(%)	N	(%)	N	(%)	N	(%)
Ceftiofur <sup>1</sup>	80	(80.8)	4	(12.5)	10	(10.1)	0	(0.0)	41	(41.4)	1	(3.1)	58	(58.6)	1	(3.1)
Tetracyclines <sup>1</sup>	31	(31.3)	0	(0.0)	18	(18.2)	0	(0.0)	15	(15.2)	0	(0.0)	24	(24.2)	1	(3.1)
Penicillin <sup>1</sup>	32	(32.3)	0	(0.0)	42	(42.4)	0	(0.0)	43	(43.4)	0	(0.0)	42	(42.4)	0	(0.0)
Ampicillin <sup>1</sup>	22	(22.2)	0	(0.0)	26	(26.3)	0	(0.0)	12	(12.1)	0	(0.0)	4	(4.0)	0	(0.0)
Sulfonamide <sup>1</sup>	20	(20.2)	0	(0.0)	10	(10.1)	0	(0.0)					4	(4.0)	0	(0.0)
Tilmicosin <sup>2</sup>	3	(3.0)	0	(0.0)												
Florfenicol <sup>2</sup>	7	(7.1)	0	(0.0)												
Amoxicillin <sup>2</sup>					15	(15.2)	0	(0.0)								
Erythromycin <sup>2</sup>					8	(8.1)	0	(0.0)								
Other	11	(11.1)	0	(0.0)	22	(22.2)	0	(0.0)	6	(6.1)						

<sup>1</sup>Antibiotic specifically mentioned in question.

<sup>2</sup>Antibiotic volunteered by producers in the "other" category.

or metritis was reported in one ORG dairy, whereas ceftiofur, tetracycline, penicillin, and ampicillin were reported for treatment of retained placenta or metritis for cows in 12 to 43% of CON herds ( $P < 0.04$ ) (Table 9). The use of systemic tetracycline and ceftiofur to treat foot problems in cows was reported in a single ORG herd compared with the use of ceftiofur, tetracycline, and penicillin to treat foot problems of cows in 24 to 59% of CON herds ( $P < 0.008$ ) (Table 9).

The use of antibiotics in footbaths (on regular schedule) to control or treat lameness was reported in 16 CON (16.2%) and three (9.4%) ORG herds ( $P = 0.34$ ), whereas the continuous use of antibiotics in footbaths was reported in 14 (14.2%) CON and one (3.1%) ORG herds ( $P = 0.09$ ). No ORG herds reported the use of tetracycline ( $P = 0.02$ ) and lincomycin ( $P = 0.06$ ) in footbaths, but these products were used in 15 (15.2%) and 11 (11.1%) CON herds, respectively. Five (5.1%) CON and three (9.4%) ORG herds utilized antibiotics other than lincomycin and tetracycline in footbaths ( $P = 0.37$ ).

There was a significant association between herd type for use of intramammary dry cow therapy ( $P < 0.001$ ). Intramammary infusion of dry cow treatment was used in 97 CON (98.0%) but only two ORG (6.3%) herds. The preparation reported by the two ORG herds was a nonantimicrobial preparation. No use of intramammary dry cow treatments was reported by 30 ORG (93.8%) and by just two (2.0%) CON farmers.

## DISCUSSION

Herds enrolled in this study were slightly larger and more productive compared with individual state averages (Table 1). The average herd size and rolling herd average reported by USDA in 2001 were 83 cows, 7794 kg (Wisconsin); 93 cows, 7950 kg (New York); 92 cows,

8765 kg (Michigan); and 67 cows, 7837 kg (Minnesota) (USDA, 2001). The larger herd size we observed is probably a result of stratified herd selection and eligibility requirements that limited enrollment of herds with <30 cows.

This study was not designed to characterize the typical ORG or CON dairy herd, but data from this study did demonstrate a number of similarities and differences based on herd type. Organic herds were smaller and produced less milk than CON herds, leading us to question the comparability of antibiotic usage based on herd type. Results of the Cochran-Mantel-Haenszel test indicated that usage of specified antibiotics was associated with herd type and was not confounded by herd size. We were unable to perform direct comparisons of other management characteristics within herd size strata because of a limited number of large ORG dairy farms enrolled in this study. It is likely that a number of the management differences we noted based on herd type were strongly related to herd size. Most CON herds housed lactating cows in free stalls and milked in a parlor, whereas most ORG herds housed and milked lactating cows in tie stalls. The maternity area was separate from the housing of lactating cows for more CON herds compared with ORG herds, probably because more ORG herds used stall barns for housing adult cows. Many larger dairy farms have had the opportunity to build modern free-stall facilities as their herds expanded, but many smaller dairy herds (both ORG and CON) are housed in older, more traditional facilities. For some smaller dairy herds, the decision to adopt an ORG production system is perceived as a way to increase the value of their product without increasing herd size.

Management differences do exist between CON and ORG herds. Several practices highly adopted by ORG herds (such as housing preweaned calves in an individ-

ual area and the use of individual water bowls for both milking and dry cows) could reduce the spread of disease. Conversely, the inability to use antimicrobials for prophylactic (dry cow therapy) and therapeutic purposes for mastitis could lead to an increase in chronic udder infections, increased SCC, and decreased milk quality on ORG farms.

Nutritional management was associated with herd type and is a possible explanation for the higher milk yield observed in CON dairy herds. Significantly more CON dairies fed cows a TMR, as well as a transition ration, and anionic salts to close-up cows. As expected, ORG herds used significantly less purchased feeds or feeds obtained from off-farm sources. The use of purchased feeds is limited for ORG dairy herds because feed must be raised organically for at least 3 yr. The only purchased feedstuffs that were used by ORG herds were whole soybeans and soybean meal. Organically raised soybeans are becoming increasingly available to support the organic farming industry.

More ORG herds used individual automatic water bowls for milking and dry cows compared with CON herds, associated with the type of housing. Shared water sources may be a potential source to spread disease among herd mates. Organic herds are required to have access to the outdoors, which probably explains the higher use of surface water for both milking cows and dry cows among ORG herds.

This study has several limitations. Most of the ORG herds were recruited by personal contact based upon proximity to universities. In most states, the enrollment process for ORG herds emulated a census, as virtually all identified ORG herds were invited to enroll. Some bias is introduced in virtually all studies that use commercial dairy herds because of the possibility that volunteer herds differ from nonresponders. The accuracy of reported antibiotic usage was not verifiable. Several dairy producers relied on memory to recall the amount and type of antibiotics, whereas others had comprehensive treatment records entered into a computer program. More CON herds recorded treatments of both lactating and nonlactating cows, but ORG herds reported significantly lower antibiotic usage. The percentage of ORG herds that did report recording treatments was similar to the percentage of herds that used antibiotics.

The potential for exposure of preweaned calves to antibiotics was higher on CON compared with ORG dairy farms. More CON dairies fed whole milk from cows that had received antibiotics and used milk replacer containing antibiotics. Similarly, few ORG dairies used coccidiostats and ionophores. These management differences are expected given the respective goals of ORG and CON herd management. Some manage-

ment characteristics of farms were similar. The quantity of colostrum fed to newborn calves and the number of herds where cattle had been diagnosed with *Salmonella* and BVD did not vary based on herd type.

During the period of this study, there was no national standard for organic food production. Whereas each certifying organization could have differences in rules for their producers, the rules appeared to be largely similar. The current ORG standards were introduced after data collection for this study was completed; however, almost all ORG herds already appeared to be in compliance with the standards before their implementation. The current ORG standards state that producers of organic livestock must not withhold medical treatment from a sick animal in order to preserve its organic status. Livestock treated with a prohibited substance are required to be identified and removed from ORG production channels. As expected, lower percentages of animals were treated with antibiotics in ORG versus CON dairies. Organic farms are allowed to use antibiotics in ill adult cows if the animal and her products (milk) are removed from organic marketing channels. No ORG herds used antibiotics to treat mastitis; however, several ORG herds did report treating respiratory disease in cows with antibiotics. When ORG dairies did use antibiotics in cows, ceftiofur was the most commonly reported drug. During the time of our study, some organic certifying organizations allowed milk to be sold from cows treated with antibiotics after an extended withdrawal period had passed. Three ORG herds did report selling ORG milk from cows that had been treated with antibiotics, and one herd did not permanently separate animals that had received antibiotics from the rest of the herd. Current organic standards do not allow use of antibiotics for use in dairy animals producing a product sold as organic.

During the period of this study, about one-third of the ORG herds reported the use of antibiotics to treat ill calves. For ORG herds that did choose to use antibiotics, penicillin was the most commonly selected antibiotic used to treat calf diarrhea and respiratory disease. The current ORG standards specify that, during the transition to ORG status, dairy animals must be under ORG management for a full year before milk production is considered organic. After a herd has been converted to ORG status, all animals must be under ORG management from the last third of gestation.

Antibiotic injection practices have been previously reported for U.S. dairy herds (National Animal Health Monitoring System, 1997). Dairy producers (n = 1219) from 20 states voluntarily responded to a questionnaire that addressed antibiotic injection practices employed during the past year. Of surveyed farms, 94 and 43% of farms reported that at least one antibiotic injection

had been administered in the past year to at least 1% and 10% of all milk cows, respectively (National Animal Health Monitoring System, 1997). Similarly, our study found that 95% of farms reported that at least one antibiotic injection was administered in the previous 2 mo to at least 1% of all milk cows. This study confirms that a very low percentage of herds do not treat cows with antibiotics. However, our study found that fewer farms (10%) reported that at least one antibiotic injection was administered in the previous 2 mo to at least 10% of all milk cows. These herds were managed conventionally.

McEwen et al. (1991) estimated usage of selected antibiotics based on mail survey results of Canadian dairy herds that had experienced antibiotic residue violations and paired control farms. Systemic antibiotics were administered to 1.3 to 1.6 cows per month (McEwen et al., 1991). Our study found that 84 of 99 CON herds treated 1 to 10% of adult cows with at least one antibiotic injection or oral dose of antibiotics in the previous 2 mo. These results are quite similar, given that the herd size of the Canadian herds was 50 (case herds) and 38 (control herds) cows. These results suggest that antibiotic usage of our study population was similar to the Canadian farm study population.

The use of intramammary dry cow treatment was highly adopted by CON dairy farms enrolled in this study. A study of 201 dairy herds in The Netherlands found that 82.8% of herds used dry cow treatment on all cows (Barkema et al., 1998). Current organic standards prohibit the use of dry cow therapy containing antibiotics, but antibiotic-free preparations are allowed.

An extensive study on drug use in Holstein calves ( $n = 104$  herds) located in Ontario was conducted between October 1980 and July 1983 (Waltner-Toews et al., 1986). Treatment days were calculated as the number of calves treated multiplied by the number of days each calf received treatment. The percent of treatment days in 4977 calves was reported for: chloramphenicol (32.0%), tetracycline (17.1%), commercial mixtures (13.7%), penicillin (13.3%), trimethoprim-sulfonamide (8.1%), sulfonamides (4.9%), and other antibiotics (10.9%). The availability and legality of antibiotics has changed since that study was conducted. The use of many antibiotics (trimethoprim-sulfa, tetracycline, penicillin, and other antibiotics) remains common for treatment of calf ailments. Several additional products (tilmicosin, florfenicol, ceftiofur, and enrofloxacin) have been developed in recent years. Enrofloxacin is approved for use in the treatment of respiratory disease in cattle. However, it is not approved for use in veal calves or cattle intended for dairy production. In the Canadian study farms, 76.9% fed milk replacer or calf starter that did not contain antibiotics, whereas 19.2% used tetracycline and 3.9% used another antibiotic in

milk replacer or calf starter (Waltner-Toews et al., 1986). Although similar drugs were used to treat calf ailments, herds enrolled in this study used considerably more antibiotics in calf starter and milk replacer compared with the Ontario dairy farms.

The results of this study show that, as expected, CON dairy farms reported significantly more use of antibiotics to treat specific ailments in both calves and cows when compared with ORG dairy farms. Most antibiotics used were approved for use in dairy cattle and appropriately used; however, a small number of herds did report the use of antibiotics that are prohibited for use in dairy. Antibiotic usage data are limited and hard to describe accurately on dairy farms. Disclosure of the data is reliant on producers' records and/or memory. In the future, more research needs to be conducted to quantify antimicrobial usage on dairy farms. As the regulations for ORG herds become more defined, further research will be needed to account for the management changes inherent in tighter regulations.

## CONCLUSION

Management differences existed between CON and ORG dairy herds in several areas. Organic herds were smaller and produced less milk compared with CON dairies. Organic dairies were more likely than CON dairies to house lactating cows in tie stalls and house preweaned calves in an individual area. More CON dairies than ORG dairies used purchased feeds and feeds obtained from off-farm sources. This study identified several characteristics of ORG herds that could reduce the need for antibiotics. However, the inability to treat mastitis and use dry cow therapy could result in higher SCC values. Reported antibiotic usage on this study was relatively low for CON dairy herds, but milk replacer containing antibiotics was used by 49.5% of enrolled CON dairy herds. Antibiotic use was slightly higher in heifer calves as compared to cows. Antibiotic use was lowest in bred heifers. Most ORG dairy farms were in compliance with current organic regulations even though the current standards had not yet been implemented.

## ACKNOWLEDGMENT

The authors thank RoseAnn Miller for technical assistance with data management.

## REFERENCES

- Barkema, H. W., Y. H. Schukken, T. J. G. M. Lam, M. L. Beiboer, G. Benedictus, and A. Brand. 1998. Management practices associated with low, medium, and high somatic cell counts in bulk milk. *J. Dairy Sci.* 81:1917-1927.

- Green, C. 2000. U. S. organic agriculture gaining ground. Economic Research Service/USDA. Agriculture Outlook. 9-14. <http://www.ers.usda.gov/epubs/pdf/agout/apr2000/ao270d.pdf>. Accessed Feb. 2003.
- Hoeben, D., C. Burvenich, and R. Heyneman. 1998. Antibiotics commonly used to treat mastitis and respiratory burst of bovine polymorphonuclear leukocytes. *J. Dairy Sci.* 81:403–410.
- McEwen, S. A., A. H. Meek, and W. D. Black. 1991. A dairy farm survey of antibiotic treatment practices, residue control methods and associations with inhibitors in milk. *J. Food Prot.* 54:454–459.
- Mitchell, J. M., M. W. Griffiths, S. A. McEwen, W. B. McNab, and A. J. Yee. 1998. Antimicrobial drug residues in milk and meat: causes, concerns, prevalence, regulations, tests, and test performance. *J. Food Prot.* 61:742–756.
- National Animal Health Monitoring System. 1997. Antibiotic injection practices on U.S. dairy operations. United States Department of Agriculture Animal and Plant Health Inspection Service. Online. Available: [http://www.aphis.usda.gov/vs/ceah/cahm/Dairy\\_Cattle/d96anti.htm](http://www.aphis.usda.gov/vs/ceah/cahm/Dairy_Cattle/d96anti.htm). Accessed February 2003.
- National Organic Program; Organic production and handling standards. October 2002. <http://www.ams.usda.gov/nop/FactSheets/ProdHandE.html>. Accessed November 6, 2002.
- SAS User's Guide: Statistics, Version 8 Edition. 1999. SAS Inst., Inc., Cary, NC.
- Sundlof, S. F., J. B. Kaneene, and R. A. Miller. 1995. National survey on veterinarian drug use in lactating dairy cows. *JAVMA* 207:347–352.
- United States Department of Agriculture's National Agricultural Statistics Service. <http://www.usda.gov/nass/>. Accessed December 5, 2002.
- Waltner-Toews, D., S. W. Martin, and A. H. Meek. 1986. Calf-related drug use on Holstein dairy farms in southwestern Ontario. *Can. Vet. J.* 27:17–22.